

FREQUENTLY ASKED QUESTIONS: Breastfeeding and COVID-19 For health care workers

(12 May 2020)



Preface


This FAQ complements the WHO interim guidance: *Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected* (13 March 2020 - [www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](http://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected)) and provides responses to questions that have arisen about the recommendations.

The interim guidance and FAQ reflect:

- i. the available evidence regarding transmission risks of COVID-19 through breastmilk;
- ii. the protective effects of breastfeeding and skin-to-skin contact, and,
- iii. the harmful effects of inappropriate use of infant formula milk.

The FAQ also draws on other WHO recommendations on Infant and Young Child Feeding and the Interagency Working Group Operational Guidance on Infant and Young Child Feeding in Emergencies. A decision tree shows how these recommendations may be implemented by health workers in maternity services and community settings, as part of daily work with mothers and families.

www.who.int/news-room/q-a-detail/q-a-on-covid-19-and-breastfeeding



1. Can COVID-19 be passed through breastfeeding?

Active COVID-19 (virus that can cause infection) has not, to date, been detected in the breastmilk of any mother with confirmed/suspected COVID-19. It appears unlikely, therefore, that COVID-19 would be transmitted through breastfeeding or by giving breastmilk that has been expressed by a mother who is confirmed/suspected to have COVID-19. Researchers continue to test breastmilk from mothers with confirmed/suspected COVID-19.

2. In communities where COVID-19 is prevalent, should mothers breastfeed?

Yes. In all socio-economic settings, breastfeeding improves survival and provides lifelong health and development advantages to newborns and infants. Breastfeeding also improves the health of mothers. In contrast, transmission of COVID-19 through breastmilk and breastfeeding has not been detected. There is no reason to avoid or stop breastfeeding.

3. Following delivery, should a baby still be immediately placed skin-to-skin and breastfed if the mother is confirmed/suspected to have COVID-19?

Yes. Immediate and continued skin-to-skin care, including kangaroo mother care, improves thermal

regulation of newborns and several other physiological outcomes, and is associated with reduced neonatal mortality. Placing the newborn close to the mother also enables early initiation of breastfeeding which also reduces neonatal mortality.

The numerous benefits of skin-to-skin contact and breastfeeding substantially outweigh the potential risks of transmission and illness associated with COVID-19.

4. If a mother is confirmed/suspected to have COVID-19, should she continue breastfeeding?

Yes. High quality evidence shows that breastfeeding reduces neonatal, infant and child mortality including in high resource settings and improves lifelong health and development in all geographies and economic settings.

The transmission of COVID-19 through breastmilk and breastfeeding has not been detected. Among the few cases of confirmed COVID-19 infection in children from other sources, most have experienced only mild or asymptomatic illness.

While breastfeeding, a mother should still implement appropriate hygiene measures, including wearing a medical mask if available, to reduce the possibility of droplets with COVID-19 being spread to her infant.

5. What are the hygiene recommendations for a breastfeeding mother confirmed/suspected to have COVID-19?

If a mother is confirmed/suspected to have COVID-19 she should:

- Wash hands frequently with soap and water or use alcohol-based hand rub, especially before touching the baby
- Wear a medical mask while feeding. It is important to:
 - Replace masks as soon as they become damp
 - Dispose of masks immediately
 - Not re-use a mask
 - Not touch the front of the mask but untie it from behind
- Sneeze or cough into a tissue, immediately dispose of it and use alcohol-based hand rub or wash hands again with soap and clean water
- Regularly clean and disinfect surfaces

6. If a mother confirmed/suspected to have COVID-19 does not have a medical face mask should she still breastfeed?

Yes. Breastfeeding unquestionably reduces neonatal and infant mortality and provides numerous lifelong health and brain development advantages to the infant/child. Mothers with symptoms of COVID-19 are advised to wear a medical mask, but even if this is not possible, breastfeeding should be continued. Other infection prevention measures, such as washing hands, cleaning surfaces, sneezing or coughing into a tissue are also important.

Non-medical masks (e.g. home-made or cloth masks) have not been evaluated. At this time, it is not possible to make a recommendation for or against their use.

7. Is it necessary for a mother with confirmed/suspected COVID-19 to wash her breast before she breastfeeds directly or before expressing milk?

If a mother is confirmed/suspected to have COVID-19 has just coughed over her exposed breast or chest, then she should gently wash the breast with soap and warm water for at least 20 seconds prior to feeding.

It is not necessary to wash the breast before every breastfeed or prior to expressing milk.

8. If a mother confirmed/suspected to have COVID-19 is not able to breastfeed what is the best way to feed her newborn/infant?

The best alternatives to breastfeeding a newborn or young infant are:

• *Expressed breastmilk*

- Expression of breastmilk is primarily done or taught through hand expression, with the use of a mechanical pump only when necessary. Hand expression and using a pump can be equally effective.
- The choice of how to express will depend on maternal preference, availability of equipment, hygiene conditions and cost.
- Expressing breastmilk is also important to sustain milk production so that mothers can breastfeed when they recover.
- The mother, and anyone helping the mother, should wash their hands before expressing breastmilk or touching any pump or bottle parts and ensure proper pump cleaning after each use. (See question 10 below)
- The expressed breastmilk should be fed to the child preferably using a clean cup and/or spoon (easier to clean), by a person who has no signs or symptoms of illness and with whom the baby feels comfortable. The mother/caregiver should wash their hands before feeding the newborn/infant.

• *Donor human milk*

- If the mother is unable to express milk and milk is available from a human milk bank, donor human milk can be fed to the baby while the mother is recovering.

• *If expressing breastmilk or donor human milk are not feasible or available, then consider:*

- Wet-nursing (another woman breastfeeds the child) (see question 11 below)
- Infant formula milk with measures to ensure that it is feasible, correctly prepared, safe and sustainable.

9. Is it safe to give expressed breastmilk from a mother confirmed/suspected to have COVID-19?

Yes. Active COVID-19 virus has not, to date, been detected in the breastmilk of any mother confirmed/suspected to have COVID-19. It is unlikely that the virus can be transmitted by giving breastmilk that has been expressed by a mother with confirmed/suspected COVID-19.

10. If a mother with confirmed/suspected COVID-19 is expressing her milk for her baby, are there extra measures needed when handling the breastmilk pump, milk storage containers or feeding utensils?

Even when COVID-19 is not a consideration, breastmilk pumps, milk storage containers and feeding utensils need to be appropriately cleaned after every use.

- Wash the pump/containers after every use with liquid soap, e.g. dishwashing liquid and warm water. Rinse after with hot water for 10-15 seconds.
- Some breast pumps parts can be put in the top rack of a dishwasher (if available). Check the instruction manual before doing this.

11. If a mother with confirmed/suspected COVID-19 is not able to breastfeed or to express breastmilk, can wet-nursing be recommended?

Wet-nursing (another woman breastfeeds the child) may be an option depending on acceptability to mothers/families, national guidelines, cultural acceptability, availability of wet-nurses and services to support mothers/wet-nurses.

- In settings where HIV is prevalent, prospective wet-nurses should undergo HIV counselling and rapid testing, according to national guidelines, where available. In the absence of testing, if feasible undertake HIV risk assessment. If HIV risk assessment/counselling is not possible, facilitate and support wet-nursing. Provide counselling on avoiding HIV infection during breastfeeding.
- Prioritise wet-nurses for the youngest infants.

12. If a mother confirmed/suspected to have COVID-19 was unable to breastfeed because she was too ill or because of another illness, when can she start to breastfeed again?

A mother can start to breastfeed when she feels well enough to do so. There is no fixed time interval to wait after confirmed/suspected COVID-19. There is no evidence that breastfeeding changes the clinical course of COVID-19 in a mother.

She should be supported in her general health and nutrition to ensure full recovery. She should also be supported to initiate breastfeeding or relactate.

13. Do the results of COVID-19 testing make any difference to infant and young child feeding recommendations?

COVID-19 testing does not have any immediate implications for decisions on infant and young child feeding.

However, confirmation of COVID-19 means that a mother should implement appropriate recommended hygiene practices for the period that she is likely to be infective i.e. while symptomatic or through the 14 days after the start of symptoms, whichever is longer.

14. Is it advisable for a mother with confirmed/suspected COVID-19 who is breastfeeding, to give a 'top-up' with infant formula milk?

No. If a mother is confirmed/suspected to have COVID-19 and is breastfeeding, there is no need to provide a 'top-up' with an infant formula milk. Giving a 'top-up' will reduce the amount of milk produced by a mother. Mothers who breastfeed should be counselled and supported to optimise positioning and attachment to ensure adequate milk production. Mothers should be counselled about responsive feeding and perceived milk insufficiency and how to respond to their infants' hunger and feeding cues to increase the frequency of breastfeeding.

15. What are key messages for a mother who wants to breastfeed but is scared about passing COVID-19 to her infant?

As part of counselling, a mother's or family's anxiety about COVID-19 should be acknowledged and responded to with the following messages:

- I. Breastfeeding and skin-to-skin contact significantly reduce the risk of death in newborns and young infants and provide immediate and lifelong health and development advantages. Breastfeeding also reduces the risk of breast and ovarian cancer for the mother.
- II. Newborns and infants are at low risk of COVID-19 infection. Among the few cases of confirmed COVID-19 infection in young children, most have experienced only mild or asymptomatic illness.
- III. The numerous benefits of breastfeeding substantially outweigh the potential risks of transmission and illness associated with COVID-19.
- IV. Active COVID-19 has not been detected in the breastmilk of any mother with confirmed/suspected COVID-19 and there is no evidence so far that the virus is transmitted through breastfeeding.

16. If a mother is confirmed/suspected to have COVID-19, is infant formula milk safer for infants?

No. There are always risks associated with giving infant formula milk to newborns and infants in all settings.

The risks associated with giving infant formula milk are increased whenever home and community conditions are compromised e.g. reduced access to health services if a baby becomes unwell / reduced access to clean water / access to supplies of infant formula milk are difficult or not guaranteed, not affordable and not sustainable.

The numerous benefits of breastfeeding substantially outweigh the potential risks of transmission and illness associated with COVID-19.

17. For what period of time are WHO recommendations on Breastfeeding and COVID-19 relevant?

The recommendations on caring and feeding of infants of mothers with confirmed/suspected COVID-19 are for the time when she is likely to be infective, i.e. while symptomatic or through the 14 days after the start of symptoms, whichever is longer.

18. Why do recommendations for mothers with confirmed/suspected COVID-19 and their infants seem different from social distancing recommendations for the general population?

Recommendations for adults and older children to maintain social distancing aim to reduce contact with asymptomatic persons who have COVID-19 and transmission of the virus that may result. This strategy will reduce the overall prevalence of COVID-19 and the number of adults who experience more serious disease.

The aim of recommendations on the care and feeding of infants and young children whose mothers have confirmed/suspected COVID-19 infection is to improve the immediate and lifelong survival, health and development of their newborns and infants. These recommendations consider the likelihood and potential risks of COVID-19 in infants and also the risks of serious illness and death when infants are not breastfed or when infant formula milk are used inappropriately as well as the protective effects of breastfeeding and skin-to-skin contact.

In general, children are at low risk of COVID-19 infection. Among the few cases of confirmed COVID-19 infection in children, most have experienced only mild or asymptomatic illness. The numerous benefits of breastfeeding substantially outweigh the potential risks of transmission and illness associated with COVID-19.

19. Is it alright for health facilities to accept free supplies of formula milk for infants of mothers with confirmed/suspected COVID-19?

No. Donations of infant formula milks should not be sought or accepted. If needed, supplies should be purchased based on assessed need. Donated formula milk is commonly of variable quality, of the wrong type, supplied disproportionate to need, labelled in the wrong language, not accompanied by an essential package of care, distributed indiscriminately, not targeted to those who need it, is not sustained, and takes excessive time and resources to reduce risks.

20. Why do WHO recommendations on mother/infant contact and breastfeeding for mothers with confirmed/suspected COVID-19 differ from those of some national and professional organizations?

WHO's recommendations on mother/infant contact and breastfeeding are based on a full consideration not only of the risks of infection of the infant with COVID-19, but also the risks of serious morbidity and mortality associated with not breastfeeding or the inappropriate use of infant formula milks as well as the protective effects of skin-to-skin contact and breastfeeding.

Recommendations of other organizations may focus only on the prevention of COVID-19 transmission without full consideration of the importance of skin-to-skin contact and breastfeeding.




www.who.int/news-room/q-a-detail/q-a-on-covid-19-and-breastfeeding

Disclaimer

The responses to questions in this document are derived from WHO publications and the Interagency Working Group Operational Guidance on Infant and Young Child Feeding in Emergencies. The WHO interim guidance was developed by a WHO global network of clinicians and clinicians who have treated patients with SARS, MERS, or severe influenza or COVID-19.

For queries, please email: outbreak@who.int with "COVID-19 clinical question" in the subject line.





DECISION TREE

for breastfeeding in context of COVID-19: Guidance for **health care and community settings**

